



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

LANE MEDICAL LIBRARY STANFORD
L293 JMS 1921 STOR
Diabetes mellitus : a system of diets /



24503420539

DIABETES MELLITUS

A SYSTEM OF DIETS


HERMAN O. MOSENTHAL, M.D.


PAUL B. HOEBER, PUBLISHER, NEW YORK

L293
JMS
1921



ANTOINE BOREL FUND





DIABETES MELLITUS

A SYSTEM OF DIETS

NOTE

The various diet tables given in this book may be obtained in pad form, 50 sheets to the pad, as follows:

- No. 1—Starch-Free Diet, Qualitative List, (page 11.)
- No. 2—Minimal Fat, Starch-Free, Measured Diet, (pages 16-17.)
- No. 3—Minimal Fat, Starch-Free, Weighed Diet, (pages 22-23.)
- No. 4—Low Fat, Starch-Free, Measured Diet, (pages 18-19.)
- No. 5—Low Fat, Starch-Free, Weighed Diet, (pages 26-27.)
- No. 6—Accessory Diet, Rich in Carbohydrates, (page 29.)

Per set of 6 pads, \$5.00; single pads, \$1.00.

Special prices on large quantities quoted on application.

PAUL B. HOEBER, *Publisher*, 67-69 East 59th Street, New York City

DIABETES MELLITUS

A SYSTEM OF DIETS

BY

HERMAN O. MOSENTHAL, M.D.

*Assistant Professor of Medicine and Attending Physician,
New York Post-Graduate Hospital and Medical School*



NEW YORK
PAUL B. HOEBER
67-69 EAST 59th STREET, NEW YORK

LARGE EDITION

COPYRIGHT, 1921
BY PAUL B. HOEBER

Published January, 1921

Printed in the United States of America

U.S. GOVERNMENT PRINTING OFFICE

PREFACE

The art of prescribing diets in diabetes mellitus has distinctly outs ability of the patient or average nurse to meet the situation. The resu that only the minority of diabetics, favored by dietitians or a specia member of the household can indulge in varied and accurate diets n grams of proteins, fats and carbohydrates. In many hospitals the s tition holds true; those having large resources have a sufficient number c to carry out the dietetic prescriptions properly while the majority rel usually very inadequate, food lists to meet the needs of the patient. A system of diets has been designed with the object of allowing any patie without special training in dietetics, to carry out the proper rationing diabetes mellitus. These diet lists have been in successful use in a nun pitals and clinics for several years. It is proposed to issue the individ pad form, so that they may be of more general use. The plan upon diets depend was first published in the *Medical Clinics of North America*; quently the scheme for the "Measured diets" was printed in *The American Journal of Medical Sciences* by Herbert Wiener and the author.




TABLE OF CONTENTS

COMMENT ON THE DIETS
DIET LISTS:
Starch-free diet, qualitative list
Measured diets:
1. Minimal fat, starch-free diet
2. Low fat, starch-free diet
Weighed diets:
1. Minimal fat, starch-free diet
2. Low fat, starch-free diet
Accessory diet, foods rich in carbohydrates
DISCUSSION OF DIETS:
Starch free diet, qualitative list
Starvation
Measured diets
Weighed diets
1. Minimal fat, starch-free diet
2. Low fat, starch-free diet
Accessory diet, foods rich in carbohydrates



COMMENT ON THE DIETS

THE diet lists virtually explain themselves. Variety in the weighed or measured diets is obtained by resorting to the vegetable or meat and fish lists. In gradations than 250 calories are desired in increasing or diminishing the this can easily be done by adding or taking away only a fraction of the foods by the diet in use differs from the next higher or lower one.

A word of explanation as to the reason for the particular proportion of protein and fats employed and as to the method of prescribing the diets may be of some value.

The minimal fat diets are intended to be emergency diets only. They are used when acidosis is to be combated. The most efficient means at hand to do prevent the accumulation of acid substances within the body, in diabetes mellitus, is to cut the fat intake to a minimum.

The so-called low fat diets are those intended for prolonged administration. Through some experimentation it was found that the lowest amount of fat, which serves to make the food palatable in the long run, is present in a starch free diet when the proteins and fats are approximately equal to each other gram for gram. At the same time the restriction of fat serves to keep the acidosis in check.

It is realized to-day that the diabetic can lose weight without detriment; what is not so generally appreciated is that this loss of weight should occur in the fat cells of the body and not in the proteins of the muscles and glands. If a depletion of protein takes place not only is weight lost, but strength and efficiency as well. In a "Sugar free" diet the only form of food which has the ability to conserve protein are the proteins themselves; fat and alcohol appear to be without value in this regard (Mosenthal, and Harrop, G. A., Jr., *Arch. Int. Med.*, 1918, XXII, 750). Hence the feeding of an excessive quantity of fat is not justified.

By clinical observation it was determined that small persons could be maintained in a condition of nitrogen (that is protein) equilibrium on the present low fat, sugar free diets of 1500 calories, larger individuals required 1750 calories (Mosenthal, and Clausen, S. W., *Arch. Int. Med.*, 1918, XXI, 269). Basing a mode of procedure upon this fact, the aim has been to have the patient acquire a sugar free urine before a diet of 1500 or 1750 calories is taken before starchy food is added. The food containing carbohydrates may be subsequently used. The table giving the foods in units of weight equivalent to one slice of bread will be found useful in carrying out this plan.

None of the proprietary foods have been advocated in the present tables. If it is desired to employ them it is wise to be guided by the list of analyses of diabetic foods as furnished by the Connecticut Agricultural Experiment Station at New Haven, Connecticut.

STARCH-FREE DIET

QUALITATIVE LIST

Many mild cases of diabetes mellitus do not require a weighed or measured diet to maintain a sugar-free urine and to control their disease. For them the starch-free diet, qualitative list, upon the opposite page will suffice. In the blank space at the bottom of this diet sheet, starch containing foods may be indicated, for use in suitable cases. The list tabulating the accessory diet of foods rich in carbohydrates will be of service in this connection.

STARCH-FREE DIET

QUALITATIVE LIST

MAY EAT:

Soups—Clear meat broths, which may contain the vegetables indicated below.
Meats—All kinds of meat, fresh, smoked or cured, except liver; all meats prepared without flour or bread-crumbs.
Fish—All kinds of fish, but no clams, oysters or scallops.
Eggs—Eggs in any form, prepared without milk, flour or sweetening (such as sugar, etc.).
Butter—Butter, oil and lard.
Cheese—All kinds of cheese.
Vegetables—Asparagus, asparagus tips, brussels sprouts, cabbage, carrots, celery, cucumbers, egg plant, endive, greens from beets, kohlrabi, leeks, pickles (sour or dill), pumpkin, radishes, rhubarb, sauerkraut, sorrel, string beans, swiss chard, tomatoes, water cress, wax beans.
Desserts—Gelatine jellies (use sour white wine, brandy or coffee for flavor).
Beverages—Tea, coffee and cocoa made from cracked cocoa (cocoa nibs), sugar, saccharine (without sugar or milk); claret, burgundy, sour white wine, whiskey in moderate amounts; carbonated waters.
Condiments—Pepper, salt, mustard, oil, vinegar, saccharine.


MUST AVOID EATING:

Sugar in any form. Bread, biscuits, and cakes of all kinds. Toast, cracker, oatmeal (and all cereals); sago, tapioca, macaroni, vermicelli, potatoes, parsnips, beets, corn, beans, peas. All fruits, fresh, preserved and dried, and jellies. Pastry, puddings and ice cream. Sauces and gravies thickened with flour.

MUST AVOID DRINKING:

Milks, ales, porter, stout, beer, cider, all sweet wines, port wine, liqueurs, wines, syrups.

BESIDES ABOVE MUST EAT:



STARVATION

When complete abstinence from food is indicated and the so-called "starvation treatment" is employed the following articles of food, whose nutritive value is negligible, are allowed: Clear broth, black coffee, plain tea, saccharine, salt and pepper. In instances the use of alcohol is permissible. Alcohol does not increase the production of sugar; it has a tendency to further the oxidation of the acid substances along the channels, thus diminishing acidosis, and it has an appreciable food value. This point to alcohol as an ideal food for the diabetic. However, its use has been largely discontinued, especially during starvation, as there are many patients in whom the patient is prone to disagree and cause nausea and vomiting. When it is desired to use a whiskey, brandy, claret or sour white wine may be ordered.

There is one point well worth bearing in mind in administering the starvation diet. This is the fact that under this form of diet the patient frequently loses a great deal of water from the tissues. This results in a feeling of weakness and lassitude which often is very marked and in advanced cases of diabetes may assume a serious character. Such a loss of fluid from the body may be prevented by the administration of considerable amounts of salt in the broth or the use of bicarbonate of soda. If a slight degree of dehydration ensues it does no harm though it should not be allowed to reach undue proportions.

THE MEASURED DIETS

The following lists have been prepared in order to bring weighed diets to their simplest possible terms. They have been designed especially for diabetic patients who are able to be up and about attending to their routine duties, and not endowed with unlimited means. The successful management of such patients depends largely upon the ease with which they are enabled to follow the prescribed dietary regime. By the use of these diet lists the physician can readily control the total caloric intake as well as the relative amounts of protein, fat and carbohydrate consumed, while at the same time the essential coöperation of the patient is reduced to a minimum in time and mental effort.

The measured diets in the modern treatment of diabetes must meet two requirements: (1) Control of the glycosuria, and (2) control of the acidosis. The first of these is accomplished by regulating the consumption of the carbohydrates and proteins, and the second by adjusting the fat intake. The greatest recent advance in the management of diabetes mellitus is the recognition of the fact that a curtailment of the fatty foods will prevent or postpone acidosis and coma. Two sets of diets are necessary, therefore: (1) In which the fats are reduced to a minimum, and (2) in which fats are allowed more liberally, but in limited amounts. The latter is the preferable diet to use when possible, as it is much the more palatable when continued for a long time. Accordingly, two diets have been designed, one the so-called "minimal fat diet" and the other the "low fat diet;" in both of them the only carbohydrate is that contained in the green vegetables.

The minimal fat diet is graded by 250 calory steps from 500 to 1250 calories. It is not feasible to increase the diet beyond this point, because when proteins make up the greater part of the nourishment, as they necessarily do in the minimal fat diet, the bulk of food becomes too great. It is even problematical whether all patients can consume the amounts designated under the 1250 calory diet. However, the safest way to curtail the food in diabetics suffering with severe acidosis or threatened coma is first to restrict the fats. By means of the present list, this can easily be done. Carbohydrates may be added if it is thought advisable.

For practical purposes of rough measurement the table and teaspoonful portions are adequate and exact within a reasonable margin of error. The portions of meat and fish are calculated to within the nearest $\frac{1}{4}$ -ounce value. One-quarter ounce is probably closer than the average scales of the butcher or home can weigh accurately, but in giving the amounts in these fractions an indication is furnished on what side allowances should be made. The actual use of these diet lists is self-explanatory.

A concrete idea of the proportion of proteins and fats in these diets may be gained from the following tables. In the minimal fat diet the ratio of protein to fat is found to be about 3 to 1 when round steak is eaten as the meat or fish, and 17 to 1 when blue fish is used. This, in either case, furnishes a very low fat intake. When the lowest possible quantity of fat is required it is evident that fish is more desirable than meat.

MEASURED DIETS

APPROXIMATE QUANTITIES OF PROTEIN, FAT AND CARBOHYDRATE IN THE MINIMAL FAT FREE DIET WHEN ROUND STEAK OR BLUE FISH ARE USED AS MEAT OR

Minimal fat diet when round steak is used:	Protein, gm.	Fat, gm.	Carbohydrate, gm.	Minimal fat diet when blue fish is used:	Protein, gm.	Fat, gm.
500 calories.....	60	20	18	500 calories.....	88	4
750 calories.....	93	33	18	750 calories.....	148	8
1000 calories.....	126	43	21	1000 calories.....	193	12
1250 calories.....	159	54	24	1250 calories.....	244	14

The next table gives the relative values of protein, fat and carbohydrate when roast beef, round steak or flounder are used as the meat or fish in the low fat diet. Particular foods were selected because they represent a high, average and low content for these diets. When roast beef is eaten the fat is higher than the protein; with the flounder the ratio is reversed. The low fat diet, when the patient chooses his food, has been calculated and found to contain, as a rule, proteins and fats in approximately equal gram for gram.

APPROXIMATE QUANTITIES OF PROTEIN, FAT AND CARBOHYDRATE IN THE LOW FAT, STEAK DIET WHEN ROAST BEEF, STEAK OR FLOUNDER ARE USED AS MEAT OR F

Low fat diet when roast beef is used:	Protein, gm.	Fat, gm.	Carbohydrate, gm.	Low fat diet when steak is used:	Protein, gm.	Fat, gm.	Carbohydrate, gm.	Low fat diet when flounder is used:	Protein, gm.
500 calories.....	22	39	12	500 calories.....	44	29	12	500 calories.....	73
750 calories.....	33	71	12	750 calories.....	72	44	12	750 calories.....	122
1000 calories.....	39	83	15	1000 calories.....	87	64	15	1000 calories.....	149
1250 calories.....	50	106	17	1250 calories.....	105	83	17	1250 calories.....	172
1500 calories.....	66	126	18	1500 calories.....	134	97	18	1500 calories.....	217
1750 calories.....	75	149	19	1750 calories.....	147	118	19	1750 calories.....	235
2000 calories.....	85	169	19	2000 calories.....	164	136	19	2000 calories.....	259

In the low fat diets there is no difficulty, as far as bulk of food is concerned, in consuming as much as 2000 calories. As Mosenthal and Clausen have shown a carbohydrate-free diet of 1500 to 1750 calories, containing approximately the same proportion of protein and fat as the present list, will maintain the nitrogen equilibrium of the patient and may be considered to furnish the patient with enough food to make him more physically efficient. If the carbohydrate tolerance of the patient permits of a still greater increase in the food it is best to add starch-containing foods and not proteins or carbohydrates.

The above food lists are presented in the belief that they will furnish a practical method of regulating the diet of sufferers from diabetes mellitus who are not receiving insulin treatment, and who, while under medical supervision, are dependent upon external resources for the details of dietetic control. These diets are not intended in any way to supplant the more accurate and ideal means of weighing and measuring food, as detailed in the subsequent diets, the weighed diets.

As printed on the following pages, the diets and their supplementary lists are given face to face. In pad form they are printed back to back, so that one sheet is a complete dietary scheme.

	Calories	500	750	1000	1250
BREAKFAST —Black coffee or plain tea as desired.....					
Meat or fish (see list) portion.....	A	4	B	C	C
Vegetables from list, heaping tablespoonsful.....			4	4	4
 DINNER —Clear meat broth, plain, as desired.....					
Meat or fish, from list, portion.....	B	4	C	D	E
Vegetables from list, heaping tablespoonsful.....			4	5	6
Pot cheese, heaping tablespoonsful.....		none	none	none	1
Gelatin jelly, flavored with coffee and saccharine, heaping tablespoonful.....		none	none	2	2
Black coffee or plain tea as desired.....					
 SUPPER —Clear meat broth, plain, as desired.....					
Meat or fish, from list, portion.....	B	4	C	D	E
Vegetables from list, heaping tablespoonsful.....			4	5	5
Black coffee or plain tea as desired.....					
 SPECIAL ORDERS —					
Saccharine, 5 tablets or less during whole day.					
No butter, fat or oil to be used in cooking.					

MINIMAL FAT STARCH-FREE DIET—MEASURED DIET

MEAT AND FISH PORTIONS						VEGETABLES (Fresh or Canned)
		Ounces of food weighed raw				
MEATS	Portion	A	B	C	D	E
Beef:						
Soup meat, fore-shank, lean.....	3	4½	7¼	9	11¾	Asparagus
Dried beef.....	1½	2½	4¼	5	6¾	Asparagus tips
Lean round steak.....	2¼	3½	5¾	7	9¼	Brussels sprouts
Lean roast beef (no visible fat).....	2¼	3½	6	7	9¼	Cabbage
Kidney.....	3	4½	7½	9	12	Cauliflower
Chicken.....	3	4½	7½	9	12	Celery
Lean leg of lamb (no visible fat).....	1¾	2½	4	5	6¾	Cucumbers
Tripe.....	5¾	8¼	14	16¾	22	Egg plant
Lean veal (no visible fat).....	2¾	4	7	8	11	Endive
FISH (Fresh only):						Greens from Kohlrabi
Bluefish.....	4	6	10	11¾	15½	Leeks
Codfish.....	4½	6½	11	13	17½	Lettuce
Flounder.....	5½	8¼	14	16½	22	Pickles, sour cream
Haddock.....	4¾	7	11¾	14¼	19	Pumpkin
Sea Bass.....	4¼	6¼	10½	12½	16½	Radishes
Sheepshead.....	3	4½	7½	9	12	Rhubarb
Smelts.....	4	6	10	11¾	15½	Sauerkraut
Weakfish.....	3¾	5½	9¼	11	14¾	Sorrel

LOW FAT, STARCH-FREE DIET—MEASURED DIET

	Calories	500	750	1000	1250	1500	1750	2000
BREAKFAST —Black coffee or plain tea as desired.								
Eggs.	1	1	1	1	2	2	2	2
Meat or fish (see list) portion.	A	B	B	C	D	E	F	F
Butter, flat teaspoonful.	none	1	1	2	2	3	3	3
DINNER —Clear meat broth, as desired.								
Meat or fish, from list, portion.	B	D	E	E	F	F	F	F
Vegetables from list, heaping tablespoonful.	4	4	5	6	6	6	6	6
Pot cheese, heaping tablespoonful.	none	none	none	none	none	1	1	1
Olive oil, teaspoonsful.	none	none	2	2	2	3	3	3
Butter, flat teaspoonsful.	1	1	2	2	2	3	3	4
Black coffee or plain tea (no sugar) as desired.
SUPPER —Clear meat broth, as desired.								
Eggs.	none	none	none	none	1	1	1	2
Meat or fish, from list, portion.	B	D	E	F	F	F	F	F
Vegetables from list, heaping tablespoonful.	4	4	5	5	6	6	6	6
Butter, flat teaspoonsful.	1	1	1	2	2	3	3	4
Black coffee or plain tea, as desired.

SPECIAL ORDERS—

Saccharine, 5 tablets or less in a day.

No extra butter, oil or fat to be used in cooking.

LOW FAT, STARCH-FREE DIET—MEASURED DIET

MEAT AND FISH PORTIONS							VEGETABLES (Fresh or Can)
	Ounces of food weighed raw						
MEATS	Portion	A	B	C	D	E	F
Bacon, fried, fat thrown away		1 1/4	1 1/2	2 3/4	3	3 3/4	5
Brain		2 1/2	3	3 1/2	6	7 1/4	9 1/2
Chicken (to be boiled or broiled)		2 1/2	3	5 1/2	6	7 1/2	10
Chicken (to be roasted)		1 1/4	1 1/2	2 3/4	2 3/4	3 1/2	4 3/4
Corned beef		1	1 1/4	2 1/4	2 1/4	2 3/4	3 3/4
Dried beef		1 1/4	1 1/2	3	3 1/4	4 1/4	5 1/2
Duck		1 1/4	1 1/4	2 1/2	2 3/4	3 1/2	4 1/2
Ham, smoked (to be boiled or broiled)		1	1 1/4	2 1/4	2 1/2	3	4
Guinea hen		2	2 1/4	4 1/4	4 1/2	5 3/4	7 1/2
Kidney		2 1/2	3	5 1/2	6	7 1/2	10
Lamb chops		0	1	1 3/4	2	2 1/2	3 1/4
Lamb (to be roasted)		1	1 1/4	2 1/4	2 1/2	3	4
Mutton (to be boiled)		1 3/4	2	4	4 1/4	5 1/4	7
Mutton chops, lean		2 1/4	2 1/2	4 3/4	5 1/4	6 1/2	8 1/2
Mutton (to be roasted)		1	1	2	2 1/4	2 3/4	3 3/4
Pork (to be roasted)		1 1/2	1 3/4	3	3 1/4	4 1/4	5 1/2
Pork chop, lean (to be broiled)		2	2 1/4	4	4 1/4	5 1/2	7
Roast beef		0	1	1 3/4	2	2 1/2	3 1/4
Soup meat, fore-shank, lean (to be boiled)		2 1/2	3	3 1/2	6	7 1/4	9 1/2
Squab		1 1/2	1 3/4	3	3 1/4	4 1/4	5 1/2
Steak, round, lean		2	2 1/4	4 1/4	4 1/2	5 3/4	7 1/2
Steak, sirloin, lean		1 3/4	2	3 3/4	4	5	6 1/2
Tongue, fresh (to be boiled)		1 3/4	2 1/4	4	4 1/4	5 1/2	7
Tongue, boiled, smoked, cold		1	1	2	2 1/4	2 3/4	3 1/2
Veal, roast or chop, lean		2 1/4	2 1/2	4 3/4	5 1/4	6 1/2	8 1/2
FISH, fresh	Portion	A	B	C	D	E	F
(To be boiled or broiled)							
Bass, black		3	3 1/2	6 1/2	7	9	11 1/2
Bass, sea		3 1/2	4 1/4	7 3/4	8 1/4	10 1/2	13 1/2
Bluefish		3 3/4	4	7 1/4	7 3/4	9 3/4	12 1/2
Butterfish		1 3/4	2	3 3/4	4	5	6 1/2
Codfish		3 3/4	4 1/4	8	8 3/4	11	14
Flounder		4 3/4	5 1/2	10 3/4	11	14	18
Haddock		4	4 3/4	8 3/4	9 1/2	12	15 1/2
Halibut		2 1/2	2 3/4	5 1/4	5 3/4	7 1/4	9 1/4
Kingfish		3 1/4	4	7 1/4	8	10	13
Mackerel		2	2 1/2	4 1/2	5	6 1/4	8
Perch		3 1/2	4 1/4	7 3/4	8 1/4	10 1/2	13 1/2
Pike		3 3/4	4 1/4	8	8 3/4	11	14
Porgy		2 1/2	2 3/4	5 1/4	5 3/4	7 1/4	9 1/4
Salmon		1 1/2	1 3/4	3	3 1/2	4 1/4	5 1/2
Sheepshead		2 1/2	3	5 1/2	6	7 1/2	9 3/4
Smelts		3 1/4	4	7 1/4	8	10	13
Weakfish		3	3 3/4	6 3/4	7 1/2	9 1/4	12
Whitefish		4	4 3/4	8 3/4	9 1/2	12	15 1/2
Canned or Smoked							
Herring, smoked		1	1 1/4	2 1/4	2 1/4	3	3 3/4
Salmon, canned		1 1/2	1 3/4	3 1/4	3 1/2	4 1/2	5 1/2
Sardines in oil		1	1 1/4	2 1/4	2 1/2	3	4
Sturgeon, smoked		1 1/4	1 1/2	3	3 1/4	4	5 1/4
Tunnyfish in oil, canned		1	1 1/4	2 1/4	2 1/2	3	4

THE WEIGHED DIETS

These diets may be employed when the means are at hand to weigh the food with some accuracy. They are preferable to the measured diets. The patient or intelligent attendant can learn to use these lists in a very short time and provide menus which present a considerable variety, please the patient's taste, as far as this is possible on a starch-free diet, and are adapted to the needs of the sufferer with diabetes mellitus.

MINIMAL FAT, STARCH-FREE DIET, WEIGHED DIET

This set of diets is intended for use in those emergencies when the lowest possible amount of fat is to be administered so as to control acidosis. When the means of weighing the food are not at hand the minimal fat, starch-free measured diet may be resorted to. These diets are not intended for long continued use. On account of the bulky character of the proteins, which make up the greater part of the ration, it is not feasible to raise these diets above 1250 grams.

A summary of the food values is as follows:

MINIMAL FAT, STARCH-FREE DIET, WEIGHED DIET

Summary of Food Values

	Calories	250	500	750	1000	1250
Protein, gm.....		33	67	111	145	183
Fat, gm.....		6	15	24	34	43
Carbohydrate, gm.....		15	18	19	23	26
Actual calories.....		253	494	556	1002	1257

If meat or fish containing less fat and more protein than the round steak be substituted, the fat content of these diets may be lowered still further.

The individual food values for some of these diets are given in the subsequent tables. A reference to these figures will enable the physician or dietitian to make finer gradations or other modifications in these diets without recourse to more elaborate tables for food values.

WEIGHED DIET
MINIMAL FAT, STARCH-FREE DIET, WEIGHED DIET
500 calories

Food		Protein, gm.	Fat, gm.	C-H, gm.	Calories
<i>Breakfast:</i>					
Steak, * round, lean, gm.....	60	16.6	4.6	0	111
Vegetables from list, gm.....	200	2.0	0	6.0	33
Black coffee or plain tea.....
<i>Dinner:</i>					
Clear meat broth, cc.....	200	4.4	0.4	0	22
Steak, * round, lean gm.....	70	19.3	5.4	0	129
Vegetables from list, gm.....	200	2.0	0	6.0	33
Black coffee or plain tea.....
<i>Supper:</i>					
Clear meat broth, cc.....	200	4.4	0.4	0	22
Steak, * round, lean gm.....	60	16.6	4.6	0	111
Vegetables from list, gm.....	200	2.0	0	6.0	33
Black coffee or plain tea.....
	...	67.3	15.4	18.0	...

MINIMAL FAT, STARCH-FREE DIET, WEIGHED DIET
1000 calories

Food		Protein, gm.	Fat, gm.	C-H, gm.	Calorie
<i>Breakfast:</i>					
Steak, * round, lean, gm.....	120	33.1	9.2	0	221
Vegetables from list, gm.....	200	2.0	0	6.0	33
Black coffee or plain tea.....
<i>Dinner:</i>					
Clear meat broth, cc.....	200	4.4	0.4	0	22
Steak, * round, lean, gm.....	150	41.4	11.6	0	278
Vegetables from list, gm.....	250	2.5	0	7.5	41
Pot cheese, gm.....	40	8.4	0.4	1.7	45
Gelatin jelly flavored with coffee and saccharine, gm.....	70	4.6	0	0	19
Black coffee or plain tea.....
<i>Supper:</i>					
Clear meat broth, cc.....	200	4.4	0.4	0	22
Steak, * round, lean, gm.....	150	41.4	11.6	0	278
Vegetables from list, gm.....	250	2.5	0	7.5	41
Black coffee or plain tea.....
	...	144.7	33.6	22.7	...

* Other meats and fish may be substituted for the round steak according to their caloric equivalents accompanying list.

On the next two pages the figures for the minimal fat, starch-free, weighed they are to be used are given. In pad form, the two pages which face each printed back to back, so that one sheet furnishes a complete dietary scheme.

MINIMAL FAT, STARCH-FREE DIET, WEIGHED DIET

	Calories	250	500	750	1000	1250
BREAKFAST —Steak, * round, lean, gm.....	20	60	100	120	160	
Vegetables from list, gm.....	100	200	200	200	200	
Black coffee or plain tea as desired.....						
 LUNCH —Clear meat broth, cc.....	200	200	200	200	250	
Steak, * round, lean, gm.....	30	70	100	150	180	
Vegetables from list, gm.....	200	200	200	250	300	
Pot cheese, gm.....	30	40	50	
Gelatin jelly flavored with coffee and saccharine, gm.	60	70	70	
Black coffee or plain tea as desired.....						
 DINNER —Clear meat broth, cc.....	200	200	200	200	250	
Steak, * round, lean, gm.....	20	60	100	150	200	
Vegetables from list, gm.....	200	200	200	250	300	
Black coffee or plain tea as desired.....						
 SPECIAL ORDERS —						
Saccharine as desired.						

* Other meats and fish may be substituted for the round steak according to their caloric equivalents from accompanying list.

MINIMAL FAT, STARCH-FREE DIET—WEIGHED DIET

CALORIC EQUIVALENT OF 10 GM. OF LEAN ROUND STEAK IN MEAT OR FISH					VEGETA Fresh or C
	Gm.	Protein, gm.	Fat, gm.	Calories	
MEATS:					
Round steak, lean.....	10	2.8	0.8	19	Asparagus
Chicken.....	11	3.5	0.5	19	Asparagus
Dried beef.....	9	3.5	0.5	19	Brussels sp
Guinea Hen.....	12	2.8	0.8	19	Cabbage
Kidney.....	17	2.8	0.8	19	Cauliflower
Mutton chop, lean.....	14	3.2	0.6	19	Celery
Mutton, boiled, lean.....	11	3.4	0.5	19	Cucumbers
Roast beef, very lean.....	17	4.0	0.3	19	Egg plant
Steak, round, lean.....	10	2.8	0.8	19	Endive
Sweet breads.....	11	4.4	0.1	19	Greens fro
Veal, roast or chop, lean.....	14	4.0	0.2	18	Kohlrabi
FISH:					
Bass, black.....	19	3.9	0.3	19	Leeks
Bass, sea.....	23	4.6	0.1	20	Lettuce
Bluefish.....	13	3.4	0.6	20	Pickles, sou
Cod fish.....	19	4.1	0.1	18	Pumpkin
Flounder.....	30	4.3	0.2	20	Radishes
Haddock.....	19	4.2	0.1	18	Rhubarb
Halibut.....	15	3.1	0.6	18	Sauerkraut
Perch.....	17	3.3	0.7	20	Sorrel
Pike.....	24	4.5	0.1	19	Spinach
Porgy.....	16	3.0	0.8	20	String bear
Shad Roe.....	15	3.1	0.6	18	Swiss charc
Smelts.....	22	3.5	0.4	18	Tomatoes
Trout.....	17	3.6	0.4	18	Water cress
Weakfish.....	20	3.6	0.5	19	Wax beans
Whitefish.....	12	2.7	0.8	19	
FISH, PRESERVED:					
Cod, salt.....	17	4.6	0.1	20	

LOW FAT, STARCH-FREE DIET—WEIGHED DIET

This series of diets is intended for routine use in those cases in which:

1. Restriction of protein is necessary to control the glycosuria.

2. There is no indication to curtail fats to the utmost because of a severe acidosis, or threatening coma.

3. The means of weighing the diet are at hand; if this is not the case the low fat, starch-free diet, measured diet, must be resorted to.

As previously stated, it requires 1500 calories of this diet series to prevent loss of protein from the body in small persons, whereas larger individuals require approximately 1750 calories to accomplish this. Foods rich in carbohydrate, according to the accessory diet list, may be added when a tolerance for the diets that prevent the loss of protein, is acquired.

A summary of the values for protein, fat, carbohydrate and calories for these diets is as follows:

	Calories	500	750	1000	1250	1500	1750	2000
Protein, gm.....	40.4	59.2	71.2	89.5	99.9	124.4	135.7	
Fat, gm.....	31.0	49.2	71.6	89.6	110.7	128.1	144.4	
Carbohydrate, gm.....	12.0	12.0	12.0	12.5	12.5	12.5	21.7	
Calories, actual.....	503	750	1006	1252	1489	1753	1986	

The individual food values for some of these diets are given. A reference to these figures will enable the physician or dietitian to make finer graduations or other modifications in these diets without recourse to more elaborate tables for food values.

500 CALORIES

Food	Gm. or c.c.	Protein, gm.	Fat, gm.	C-H, gm.	Calories	Calories per meal
<i>Breakfast:</i>						
One egg	50	6.6	6.0	0	83	
Bacon	40	4.2	7.6	0	88	
Black coffee	171
<i>Dinner:</i>						
Broth	150	3.3	0.3	.0	16	
Steak ¹	40	9.4	4.1	0	77	
Vegetables ²	200	2.0	0	6.0	33	
Butter	5	0.1	4.3	0	40	
Black coffee	166
<i>Supper:</i>						
Broth	150	3.3	0.3	0	16	
Steak ¹	40	9.4	4.1	0	77	
Vegetables ²	200	2.0	0	6.0	33	
Butter	5	0.1	4.3	0	40	
Plain tea	166
		40.4	31.0	12.0	..	503

¹ The bacon is weighed uncooked. The fat and protein is calculated for the cooked product.

² The caloric equivalent of other meat or fish should be frequently substituted from the list on page 27 to furnish variety in the diet.

³ Two or three different vegetables should be chosen from the list on page 27 which tabulates the vegetables containing 5 per cent. or less of carbohydrates.

1000 CALORIES

Food	Gm. or c.c.	Protein, gm.	Fat, gm.	C-H, gm.	Calories	C per
<i>Breakfast:</i>						
Eggs (2).....	100	13.2	12.0	0	166	
Bacon (see foot note 1, page 24).....	50	5.3	9.6	0	111	
Butter.....	5	0.1	4.3	0	40	
Black coffee.....	
<i>Dinner:</i>						
Broth.....	150	3.3	0.3	0	16	
Steak (see foot note 2, page 24).....	100	23.9	10.2	0	193	
Vegetables (see foot note 3, page 24).....	200	2.0	0	6.0	33	
Olive oil.....	10	0	10.0	0	93	
Butter.....	10	0.1	8.6	0	80	
Black coffee.....	
<i>Supper:</i>						
Broth.....	150	3.3	0.3	0	16	
Steak (see foot note 2, page 24).....	75	17.9	7.7	0	145	
Vegetables (see foot note 3, page 24).....	200	2.0	0	6.0	33	
Butter.....	10	0.1	8.6	0	80	
Tea (plain).....	
		71.2	71.6	12.0	

1500 CALORIES

Food	Gm. or c.c.	Protein, gm.	Fat, gm.	C-H, gm.	Calories	C per
<i>Breakfast:</i>						
Eggs (2).....	100	13.2	12.0	0	166	
Bacon (see foot note 1, page 24).....	60	6.4	11.5	0	133	
Butter.....	10	0.1	8.6	0	80	
Black coffee.....	
<i>Dinner:</i>						
Broth.....	150	3.3	0.3	0	16	
Steak (see foot note 2, page 24).....	140	33.5	14.3	0	270	
Vegetables (see foot note 3, page 24).....	200	2.0	0	6.0	33	
Cream cheese.....	20	5.2	6.7	0.5	86	
Olive oil.....	15	0	15.0	0	140	
Butter.....	15	0.2	12.9	0	120	
Black coffee.....	
<i>Supper:</i>						
Broth.....	150	3.3	0.3	0	16	
One egg.....	50	6.6	6.0	0	83	
Steak (see foot note 2, page 24).....	100	23.9	10.2	0	193	
Vegetables (see foot note 3, page 24).....	200	2.0	0	6.0	33	
Butter.....	15	0.2	12.9	0	120	
Tea (plain).....	
		99.9	110.7	12.5	

2000 CALORIES

Food	Gm. or c.c.	Protein, gm.	Fat, gm.	C-H, gm.	Calories	C per
<i>Breakfast:</i>						
Eggs (2).....	100	13.2	12.0	0	166	
Ham.....	75	15.2	16.8	0	219	
Butter.....	15	0.2	12.9	0	120	
Vegetables (see foot note 3, page 24).....	100	1.0	0	3.0	16	
Black coffee.....	
<i>Dinner:</i>						
Broth.....	160	3.5	0.3	0	17	
Steak (see foot note 2, page 24).....	160	38.2	16.3	0	308	
Vegetables (see foot note 3, page 24).....	300	3.0	0	9.0	49	
Cream cheese.....	30	7.8	10.1	0.7	129	
Butter.....	20	0.2	17.2	0	160	
Olive oil.....	15	0	15.0	0	140	
Black coffee.....	
<i>Supper:</i>						
Broth.....	160	3.5	0.3	0	17	
Eggs (2).....	100	13.2	12.0	0	166	
Steak (see foot note 2, page 24).....	140	33.5	14.3	0	270	
Vegetables (see foot note 3, page 24).....	300	3.0	0	9.0	49	
Butter.....	20	0.2	17.2	0	160	
Tea (plain).....	
		135.7	144.4	21.7	

On the next two pages the figures for the low fat, starch-free, weighed diets as are to be used are given. In pad form, the two pages which face each other, are printed back to back, so that one sheet furnishes a complete dietary scheme.

LOW FAT, STARCH-FREE DIET, WEIGHED DIET

Calories	500	750	1000	1250	1500	1750	2000
BREAKFAST —Eggs, gm.....	(1)50	(1)50	(2)100	(2)100	(2)100	(2)100	(2)100
Bacon, * gm.....	40	40	50	60	60	75	75
Ham, gm.....	10	15
Butter, gm.....	5	5	10	...	100
Vegetables, ** gm.....
Black coffee.....							
 DINNER —Broth, cc.....	150	150	150	150	150	150	160
Steak (Sirloin), *** gm.....	40	90	100	125	140	160	160
Vegetables, ** gm.....	200	200	200	200	200	200	300
Cream cheese, gm.....	20	20	25	30
Butter, gm.....	5	5	10	15	15	15	20
Olive oil, cc.....	...	10	10	10	15	15	15
Black coffee.....							
 SUPPER —Broth, cc.....	150	150	150	150	150	150	160
Eggs, gm.....	(1)50	(1)50	(2)100
Steak (Sirloin), *** gm.....	40	70	75	100	100	140	140
Vegetables, ** gm.....	200	200	200	200	200	200	300
Butter, gm.....	5	5	10	10	15	20	20
Tea (plain)							
 SPECIAL ORDERS —							
Saccharine as desired.							

* The bacon is weighed uncooked. The fat and protein content is calculated for the cooked product.

** Two or three different vegetables should be chosen from the accompanying list which tabulates the vegetables containing 5 per cent. or less of carbohydrates.

*** The caloric equivalent of other carbohydrate-free meat or fish should be frequently substituted from the accompanying list to furnish variety in the diet.

LOW-FAT, STARCH-FREE DIET—WEIGHED DIET

CALORIC EQUIVALENT OF 10 GM. OF SIRLOIN STEAK IN MEAT OR FISH					VEGETABLES (Fresh or Canned)
Food	Gm.	Protein, gm.	Fat, gm.	Calories	
<i>Meats:</i>					
Sirloin steak.....	10	2.4	1.0	19	Asparagus
Bacon, fried, fat discarded.....	7	0.9	1.6	19	Asparagus
Brains, beef.....	16	1.4	1.5	19	Brussels sprouts
Capon.....	9	2.4	1.0	19	Cabbage
Chicken.....	11	3.5	0.5	19	Cauliflower
Corned beef.....	6	0.9	1.6	19	Celery
Dried beef.....	9	3.5	0.5	19	Cucumbers
Duck.....	6	1.0	1.6	19	Egg plant
Ham, fresh.....	5	0.8	1.7	19	Endive
Ham, smoked.....	4	0.7	1.6	18	Greens (fresh)
Ham, smoked, boiled.....	7	1.4	1.6	20	Kohlrabi
Ham, smoked, fried.....	5	1.1	1.7	20	Leeks
Guinea hen.....	12	2.8	0.8	19	Lettuce
Kidney.....	17	2.8	0.8	19	Pickles, sour
Lamb chop.....	5	1.1	1.5	19	Pumpkin
Lamb, roast.....	10	2.0	1.3	20	Radishes
Mutton chop, lean.....	14	3.2	0.6	19	Rhubarb
Mutton, boiled, lean.....	11	3.4	0.5	19	Sauerkraut
Mutton, roast.....	6	1.5	1.4	19	Sorrel
Pork chop, lean.....	12	3.1	0.7	19	Spinach
Pork, roast.....	9	2.6	0.9	19	String beans
Roast beef.....	5	1.1	1.4	18	Swiss chard
Roast beef, very lean.....	17	4.0	0.3	19	Tomatoes
Squab.....	7	1.3	1.6	20	Watercress
Steak round, lean.....	10	2.8	0.8		
Steak, sirloin, lean.....	10	2.4	1.0		
Steak, tenderloin.....	7	1.7	1.4		
Sweet breads.....	11	4.4	0.1		
Tongue beef.....	12	2.3	1.1		
Turkey, roast.....	7	2.0	1.3		
Veal, roast or chop, lean.....	14	4.0	0.2		
<i>Fish, fresh:</i>					
Bass, black.....	19	3.9	0.3	19	
Bass, sea.....	23	4.6	0.1	20	
Bluefish.....	13	3.4	0.6	20	
Butterfish.....	11	2.0	1.2	19	
Codfish.....	19	4.1	0.1	18	
Flounder.....	30	4.3	0.2	20	
Haddock.....	19	4.2	0.1	18	
Halibut.....	15	3.1	0.6	18	
Mackerel.....	14	2.4	1.0	19	
Perch.....	17	3.3	0.7	20	
Pike.....	24	4.5	0.1	19	
Porgy.....	16	3.0	0.8	20	
Salmon.....	11	2.2	1.1	19	
Shad.....	12	2.3	1.1	20	
Shad roe.....	15	3.1	0.6	18	
Smelts.....	22	3.5	0.4	18	
Trout.....	17	3.6	0.4	18	
Weakfish.....	20	3.6	0.5	19	
White fish.....	12	2.7	0.8	19	
<i>Fish, preserved:</i>					
Cod, salt.....	17	4.6	0.1	20	
Herring, smoked.....	6	2.2	1.0	18	
Mackerel, salt.....	6	1.0	1.6	19	
Salmon, canned.....	10	2.2	1.2	20	
Sardines, canned.....	7	1.6	1.4	20	


THE ACCESSORY DIET OF FOODS RICH IN CARBOHYDRATES

If the patient's urine continues to be sugar free on a "carbohydrate-free" diet of sufficient caloric value (1500 to 1750 calories), carbohydrate-containing foods may be added and the carbohydrate tolerance of the patient be determined. In those cases able to utilize a considerable amount of starch, the accessory diet may be varied from day to day, and use may be made of the following table, which gives the carbohydrate equivalent of one slice (1 ounce or 30 grams) of white bread, containing approximately 5 grams of starch.


THE ACCESSORY DIET OF FOODS RICH IN CARBOHYDRATE


EACH PORTION CONTAINS APPROXIMATELY 15 GRAMS OF CARBOHYDRATES

Foods	Household Measure	Gm.	Foods	Household Measure
<i>Uncooked Flours, etc.:</i>				
Barley.....	1 h. tbsp.	21	Apple.....	1 med
Buckwheat.....	1 h. tbsp.	19	Apricots.....	2 larg
Cornmeal.....	1 h. tbsp.	20	Banana (without skin).....	½ med
Farina.....	1 h. tbsp.	20	Cherries.....
Hominy.....	1 h. tbsp.	18	Currants.....	5 h. tl
Macaroni.....	1 h. tbsp.	20	Grapefruit.....	½ sma
Noodles.....	1½ h. tbsp.	20	Huckleberries.....	3½ h. tl
Oatmeal.....	1 h. tbsp.	22	Lemons.....	2 med
Rice.....	1 h. tbsp.	18	Muskmelon.....	½
Rye flour.....	1 h. tbsp.	18	Nectarine.....	1
Spaghetti.....	1½ h. tbsp.	20	Olives, green.....	20
Vermicelli.....	1½ h. tbsp.	21	Orange.....	½ larg
Wheat flour.....	1 h. tbsp.	20	Peaches.....	1½ med
<i>Bread and Crackers:</i>				
Bread.....	1 slice.	30	Pear.....	1 sma
Breakfast biscuit (Huntley & Palmer).....	3	18	Pineapple.....	3 slice
Cornbread.....	1 slice.	32	Plums.....	2 med
Roll (Vienna).....	½	25	Raspberries.....	4½ h. tl
Uneeda Biscuit.....	3	18	Strawberries.....	8 h. tl
Zwieback.....	1½	20	Watermelon.....	Large slice
<i>Cooked Cereals:</i>				
“Force”.....	5 h. tbsp.	18	Dried Fruits:	
Farina.....	2½ h. tbsp.	125	Apples.....	3 sma
Grapenuts.....	1½ h. tbsp.	20	Apricots.....	3 larg
Hominy.....	1½ h. tbsp.	90	Currants.....	1½ tbsp
Macaroni.....	2 h. tbsp.	100	Dates.....	3
Oatmeal.....	2½ h. tbsp.	130	Figs.....	1 larg
Rice.....	1½ h. tbsp.	60	Prunes.....	2 larg
Shredded wheat biscuit.....	¾	22	Raisins.....	10 larg
<i>Cooked Vegetables:</i>				
Artichokes.....	1 medium.	320	Milk and Cream:	
Beans (baked—canned).....	2 h. tbsp.	75	Buttermilk.....	1½ tum
Beans, lima.....	1¼ h. tbsp.	50	Cream, 16 per cent.....	1½ tum
Beets.....	6 h. tbsp.	200	Cream, 40 per cent.....	1½ tum
Carrots.....	13 h. tbsp.	440	Koumiss.....	1½ tum
Okra.....	4 h. tbsp.	200	Whole milk.....	1½ tum
Onions.....	3	300	Nuts:	
Parsnips.....	4 slices.	120	Almonds.....	60
Peas, green.....	3 h. tbsp.	100	Brazil.....	30
Potato (baked).....	½ medium.	60	Chestnuts (roasted).....	15
Potato (boiled).....	½ medium.	70	Cocoanut.....	1 slice (3)
Potato (mashed).....	1½ h. tbsp.	80	Filberts.....	100
Potato, sweet (boiled).....	½ medium.	35	Peanuts.....	40
Squash.....	2 h. tbsp.	100	Pecans.....	35
Turnips.....	3 h. tbsp.	210	Pistachio.....	190
			Walnuts.....	30









LANE MEDICAL LIBRARY

To avoid fine, this book should be returned on
or before the date last stamped below.

--	--	--	--

L793 Monenthal, H.O.
M89 Diabetes mellitus.
1971 53422

5 X 801

DATE DUE

